



Total Syntheses of Conidiogenone B, Conidiogenone, Conidiogenone C, Conidiogenone D, and Conidiogenol

Po Yuan 05.05.21

P. Hu, H. M. Chi, K. C. DeBacker, X. Gong, J. H. Keim, I. T. Hsu, S. A. Snyder, *Nature* **2019**, 569, 703 – 707.



Synthesis of Hoveyda's NHC•Ag



Step 2: Enantioselective Synthesis of All-Carbon Quaternary Stereogenic Centers: *Angew. Chem. Int. Ed.* **2008**, *47*, 7358 – 7362.

Step 5: Baran's reductive coupling: *J. Am. Chem. Soc.* **2014**, *136*, 1304–1307.

Step 6 and 7: Corey's hydrazone-mediated alkylation: *Tetrahedron Lett.* **1976**, *17*, 3–6.

Step 11: A radical-mediated, chemoselective NHK-type reaction: Recent advances in the asymmetric Nozaki–Hiyama–Kishi reaction. *Synthesis* **2016**, *48*, 4038–4049.

Toolbox Approach to the Search for Effective Ligands for Catalytic Asymmetric Cr-Mediated Coupling Reactions. *J. Am. Chem. Soc.* **2009**, *131*, 15387–15393.